Serial No.: 09/889,645 Filed: January 24, 2002

Page 2 of 8

Listing of Claims

The following list of claims will replace all prior versions and listings of claims in the application.

- 1. (Currently Amended) A method of removal of abnormal infective prion proteins associated with transmissible spongiform encephalopies (TSEs) from an aqueous liquid containing a natural product, which consists essentially of passing the liquid through a depth filter formed of a matrix comprising (a) a binder and (b) kieselguhr or perlite particles or mixtures thereof and having a pore size providing a retention less than 6 μ m, and so removing abnormal infective prion proteins which may be present in the liquid such that the liquid is non-infective with respect to prion protein infectivity, wherein the depth filter is a single use filter aqueous liquid is a blood plasma product derived from plasma.
 - 2. (Canceled)
- 3. (Previously Presented) The method according to claim 1, wherein the binder is cellulose.
 - 4-5. (Canceled)
- 6. (Previously Presented) The method according to claim 1, carried out in the absence of cationic or anionic charged material.
- 7. (Previously Presented) The method according to claim 1 carried out at a pH in the range 4 to 10.
- 8. (Previously Presented) The method according to claim 1, wherein the pore size is in the range 0.6 to 6 microns.
 - 9. (Previously Presented) The method according to claim 1, wherein the pore

Serial No.: 09/889,645 Filed: January 24, 2002

Page 3 of 8

size is in the range 0.6 to 1.5 microns.

10. (Previously Presented) The method according to claim 1, wherein the depth filter has a thickness of 2 to 5 mm.

11. (Canceled)

- 12. (Currently Amended) The method according to claim 1, wherein the aqueous liquid comprises a blood plasma product derived from human plasma is human plasma.
- 13. (Previously Presented) The method according to claim 12, wherein the blood plasma product is selected from the group consisting of albumin, an immunoglobulin, Factor IX, thrombin, fibrinogen, Factor VIII, Factor II, Factor VIII, Factor IX, and Factor X.
- 14. (Previously Presented) A liquid subjected to prion removal according to the method of claim 1.
- 15. (Currently Amended) The method according to claim 1, wherein the aqueous liquid comprises an animal-derived <u>a</u> product selected from the group consisting of heparin and hormones.
- 16. (Previously Presented) The method according to claim 1, wherein the abnormal infective prion protein is associated with conditions selected from the group consisting of Creutzfeldt-Jakob Disease, variant Creutzfeldt-Jakob Disease, bovine spongiform encephalopy and scrapie.

17-24. (Canceled)

Serial No.: 09/889,645 Filed: January 24, 2002

Page 4 of 8

25. (Currently Amended) The method according to claim $\underline{1}$ [[12]], wherein the blood plasma product is selected from the group consisting of immunoglobulins and albumin.

26-27. (Canceled)

28. (Previously Presented) The method of claim 1, wherein the filter is pretreated with ethanol.

29-30. (Canceled)

- 31. (Currently Amended) A method of removal of abnormal infective prion proteins associated with transmissible spongiform encephalopies (TSEs) from an aqueous liquid containing a natural product a plasma fraction, which wherein the method consists essentially of passing the liquid plasma fraction through a depth filter pretreated with ethanol and formed of a matrix comprising (a) a binder and (b) kieselguhr or perlite particles or mixtures thereof and having a pore size providing a retention less than 6 μ m, and so removing abnormal infective prion proteins which may be present in the liquid such that the liquid is non-infective with respect to prion protein infectivity.
- 32. (Previously Presented) The method of claim 1, wherein the depth filter has a permeability of 110 or 220 L/m²/min.
 - 33. (New) The method of claim 1, wherein the depth filter is a single use filter.
- 34. (New) The method of claim 1, wherein the aqueous liquid is a cell-free blood plasma product.
- 35. (New) The method of claim 31, wherein the plasma fraction consists essentially of a protein selected from the group consisting of immunoglobulins and albumin.

Serial No.: 09/889,645 Filed: January 24, 2002

Page 5 of 8

36. (New) The method of claim 31, wherein the depth filter is pretreated with ethanol.

37. (New) The method of claim 31, wherein the depth filter is a single use filter.